

Shear viscosity of the gluon plasma in the stochastic-vacuum approach

Dmitri Antonov

Fakultät für Physik, Universität Bielefeld, D-33501 Bielefeld, Germany

Shear viscosity of the gluon plasma in SU(3) YM theory is calculated nonperturbatively, within the stochastic vacuum model. The result for the ratio of the shear viscosity to the entropy density, proportional to the squared chromo-magnetic gluon condensate and the fifth power of the correlation length of the chromo-magnetic vacuum, falls off with the increase of temperature. At temperatures larger than the deconfinement critical temperature by a factor of 2, this fall-off is determined by the sixth power of the temperature-dependent strong-coupling constant and yields an asymptotic approach to the conjectured lower bound of $1/(4\pi)$, achievable in $\mathcal{N} = 4$ SYM theory. As a by-product of the calculation, we find a particular form of the two-point correlation function of gluonic field strengths, which is the only one consistent with the Lorentzian shape of the shear-viscosity spectral function.

I. INTRODUCTION AND PRELIMINARY ESTIMATES

The RHIC data on collective expansion dynamics of the quark-gluon plasma can be described by the relativistic hydrodynamics applied to a system with very large initial pressure gradients [1]. According to these data, particles of different mass are emitted from the quark-gluon-plasma fireball with a common fluid velocity, that is a signature of a hydrodynamic expansion. Due to a large elliptic flow in noncentral collisions [2], an agreement between the experimental data [2, 3] and the predictions of relativistic hydrodynamics can only be reached when the flow of the QGP-fluid is treated as almost non-viscous. This leads to an indication that, in the vicinity of the deconfinement phase transition, the quark-gluon plasma produced in the RHIC experiments behaves more like an ideal quantum liquid rather than a weakly interacting gas. The mean free path L_{mfp} of a parton, which traverses such a liquid, is much smaller than the inter-particle distance, which is of the order of the inverse temperature $\beta = 1/T$, i.e. $(L_{\text{mfp}}^{\text{liq}}/\beta) \ll 1$.

One can consider for comparison a weakly interacting dilute-gas model of the quark-gluon plasma. There $L_{\text{mfp}}^{\text{gas}} \sim (\rho\sigma_t)^{-1}$ with ρ and σ_t standing for the particle-number density and the Coulomb transport cross section, respectively. Using the standard estimates $\rho \sim T^3$ and $\sigma_t \sim g^4\beta^2 \ln g^{-1}$, where $g = g(T)$ is the perturbative finite-temperature QCD coupling, one obtains $(L_{\text{mfp}}^{\text{gas}}/\beta) \sim 1/(g^4 \ln g^{-1}) \gg 1$, that strongly contradicts the above-mentioned experimental results. One can check [4] that these results could have only been reproduced by the dilute-gas model if the perturbative transport cross section, σ_t , were larger by an order of magnitude. This inconsistency of the weakly interacting quark-gluon plasma with the RHIC data initiated recent calculations of kinetic coefficients in the *strongly* interacting relativistic plasmas.

Among these coefficients, the one whose values define whether the plasma can be considered as weakly or strongly interacting is the shear viscosity η . It is related to the above (L_{mfp}/β) -ratio via the estimate $(\eta/s) \sim (L_{\text{mfp}}/\beta)$, where s is the entropy density. According to this relation, the shear-viscosity to the entropy-density ratio, η/s , becomes smaller when the plasma interacts stronger. For instance, for $T \sim 200$ MeV and the estimated typical mean free path $L_{\text{mfp}} \sim 0.1$ fm, one has $(\eta/s) \sim 0.1$. On the other hand, since the mean momentum change Δp of a parton, which propagates through the plasma over the distance L_{mfp} , is of the order of T , the Heisenberg uncertainty principle forbids the ratio $(L_{\text{mfp}}/\beta) \sim L_{\text{mfp}} \cdot \Delta p$ (and therefore also η/s) to be vanishingly small. Up to now, the minimal value of $1/(4\pi) \simeq 0.08$ for the shear-viscosity to the entropy-density ratio has been found in $\mathcal{N} = 4$ SYM theory [5]. It is thus challenging to find other QCD-motivated models where this ratio would be that small. Recently, it has been demonstrated [6] that such low values of the (η/s) -ratio can take place even in the perturbative YM plasma, due to the bremsstrahlung $gg \leftrightarrow ggg$ processes. In Ref. [7] it has been argued, though, that the perturbatively calculated collisional viscosity is anyhow larger (and therefore subdominant) compared to the so-called anomalous viscosity, which is generated by plasma instabilities.

In this paper, we calculate the (η/s) -ratio in the gluon plasma of SU(3) YM theory nonperturbatively. We obtain the shear viscosity from the Kubo formula, which relates the corresponding spectral density $\rho(\omega)$ to the Euclidean correlation function of the $(1,2)$ -component of the energy-momentum tensor $T_{12}(\mathbf{x}, x_4)$. This method, proposed in Ref. [8], has been explored in Refs. [9, 10] with the aim to simulate shear viscosity on the lattice. Here we work in the continuum limit and parametrize the Euclidean correlation function of

the energy-momentum tensors by means of the stochastic vacuum model [11]. This model generalizes QCD sum rules by assuming the existence of not only the gluon condensate $\langle g^2(F_{\mu\nu}^a)^2 \rangle$ but also of the finite vacuum correlation length μ^{-1} . This assumption is justified by the lattice results on the exponential fall-off at large distances of the two-point correlation function of gluonic field strengths [13, 14], $\langle F_{\mu\nu}^a(x) F_{\lambda\rho}^b(0) \rangle \sim e^{-\mu|x|}$. By virtue of this finding, the model manages to quantitatively describe confinement; for instance, the string tension reads $\sigma \propto \mu^{-2} \langle g^2(F_{\mu\nu}^a)^2 \rangle$.

Below we will use a finite-temperature generalization of the stochastic vacuum model, accessible by implementing the Euclidean periodicity of the x_4 -coordinate. In the deconfinement phase of interest, such a generalization yields for the spatial string tension $\sigma_s(T)$ a formula [15] pretty similar to its above-quoted vacuum counterpart. This formula reads $\sigma_s(T) \propto \mu^{-2}(T) \langle g^2(F_{ij}^a)^2 \rangle_T$, where $\mu^{-1}(T)$ is the correlation length of the chromo-magnetic vacuum, and $\langle g^2(F_{ij}^a)^2 \rangle_T$ is the chromo-magnetic gluon condensate, which survives the deconfinement phase transition. The temperature dependence of the two main ingredients of the model, $\mu(T)$ and $\langle g^2(F_{ij}^a)^2 \rangle_T$, can be extracted from the results of the lattice simulations [13, 16].

Since $T_{12} = g^2 F_{1\mu}^a F_{2\mu}^a$, one *a priori* expects from the Kubo formula, where the $\langle T_{12}(0) T_{12}(x) \rangle$ -correlator enters, that the shear viscosity $\eta \propto \langle g^2(F_{ij}^a)^2 \rangle_T^2$. This is a general prediction of the stochastic vacuum model for all the kinetic coefficients, for example for the jet quenching parameter [17]. In fact, according to the Kubo formula, all the kinetic coefficients are proportional to the total scattering cross section of the propagating parton, which itself is proportional to $\langle g^2(F_{ij}^a)^2 \rangle_T^2$ in this model [18, 19]. Therefore, since the shear viscosity has the dimensionality of [mass]³, one can on entirely dimensional grounds expect for it the following result:

$$\eta \propto \mu^{-5}(T) \langle g^2(F_{ij}^a)^2 \rangle_T^2.$$

At temperatures larger than the temperature of dimensional reduction, $T > T_*$, $\mu(T)$ and $\langle g^2(F_{ij}^a)^2 \rangle_T^2$ are proportional to the corresponding power of the only dimensionful parameter present in the YM action at such temperatures, $g^2 T$, i.e.

$$\mu(T) \propto g^2 T, \quad \langle g^2(F_{ij}^a)^2 \rangle_T \propto (g^2 T)^4.$$

On the other hand, the entropy density $s(T) \propto T^3$, so that

$$\frac{\eta}{s} \propto g^6(T) \quad \text{at} \quad T > T_*.$$

In this paper, we quantitatively answer the naturally arising question of whether or not this function manages to get below the $(4\pi)^{-1}$ -threshold at temperatures $T \lesssim 5T_c$, which are accessible experimentally and on the lattice.

The outline of the paper is as follows. In Section II, by assuming an exponential fall-off for the two-point correlation function of the energy-momentum tensors $\langle T_{12}(0)T_{12}(x) \rangle$, we obtain from the Kubo formula an integral equation for the spectral density $\rho(\omega)$ of η . In Section III, by using for $\rho(\omega)$ a Lorentzian-type *ansatz*, with the same correlation length as that of $\langle T_{12}(0)T_{12}(x) \rangle$, we explore this equation for the cases of large and small $|k|$'s, where k is the number of a Matsubara mode. The solution in the large- $|k|$ limit yields the range of variation of the numerical parameter α , which enters the initial parametrization of $\langle T_{12}(0)T_{12}(x) \rangle$. The solution in the small- $|k|$ limit can only coincide with the large- $|k|$ solution for a single value of α from this range. This fixes α completely and makes further calculations straightforward. In Section IV, we analytically calculate the shear viscosity η . In Section V, we use this result for η to numerically find the ratio η/s . Also in Section V we compare the calculated nonperturbative spectral density $\rho(\omega)$ with the perturbative one, which dominates at large ω 's. In Section VI, we discuss the results of the paper, as well as possible further developments. In Appendix A, we illustrate the separation of perturbative contributions to the Kubo formula from the nonperturbative ones.

II. KUBO FORMULA FOR THE SPECTRAL DENSITY

Shear viscosity η can be defined through the relation

$$\eta = \pi \left. \frac{d\rho}{d\omega} \right|_{\omega=0}, \quad (1)$$

where the spectral density $\rho(\omega)$ is a solution to the following integral equation, called Kubo formula [8, 10]

$$\int_0^\infty d\omega \rho(\omega) \frac{\cosh \left[\omega \left(x_4 - \frac{\beta}{2} \right) \right]}{\sinh(\omega\beta/2)} = \int d^3x \sum_{n=-\infty}^{+\infty} \langle T_{12}(0)T_{12}(\mathbf{x}, x_4 - \beta n) \rangle. \quad (2)$$

Here the sum on the RHS runs over winding modes. We use the Fourier decomposition on the LHS of Eq. (2):

$$\frac{\cosh \left[\omega \left(x_4 - \frac{\beta}{2} \right) \right]}{\sinh(\omega\beta/2)} = 2T \cdot \omega \sum_{k=-\infty}^{+\infty} \frac{e^{i\omega_k x_4}}{\omega^2 + \omega_k^2}, \quad (3)$$

where $\omega_k = 2\pi T k$ is the k -th Matsubara frequency. The idea is to have a similar decomposition also on the RHS of Eq. (2). For the implementation of this idea, the following chain of equalities is important:

$$\begin{aligned}
\sum_{k=-\infty}^{+\infty} \frac{e^{i\omega_k x_4}}{[1 + (\omega_k/M)^2]^\alpha} &= \frac{1}{\Gamma(\alpha)} \sum_{k=-\infty}^{+\infty} \int_0^\infty d\lambda \lambda^{\alpha-1} e^{-\lambda[1 + (\omega_k/M)^2] + i\omega_k x_4} = \\
&= \frac{M\beta}{2\sqrt{\pi}\Gamma(\alpha)} \int_0^\infty d\lambda \lambda^{\alpha-\frac{3}{2}} e^{-\lambda} \sum_{n=-\infty}^{+\infty} e^{-\frac{M^2(x_4 - \beta n)^2}{4\lambda}} = \\
&= \frac{M^4\beta}{16\pi^2\Gamma(\alpha)} \int_0^\infty d\lambda \lambda^{\alpha-3} e^{-\lambda} \int d^3x \sum_{n=-\infty}^{+\infty} e^{-\frac{M^2[\mathbf{x}^2 + (x_4 - \beta n)^2]}{4\lambda}} = \\
&= \frac{M^4\beta}{2^{\alpha+1}\pi^2\Gamma(\alpha)} \int d^3x \sum_{n=-\infty}^{+\infty} \frac{K_{2-\alpha}\left(M\sqrt{\mathbf{x}^2 + (x_4 - \beta n)^2}\right)}{\left[M\sqrt{\mathbf{x}^2 + (x_4 - \beta n)^2}\right]^{2-\alpha}}.
\end{aligned}$$

Here $\alpha > 0$ is some parameter, “ Γ ” and “ $K_{2-\alpha}$ ” stand for the Gamma and the MacDonald functions, respectively. We assume that, at $T = 0$,

$$\langle T_{12}(0)T_{12}(x) \rangle = N(\alpha) \langle G^2 \rangle^2 \frac{K_{2-\alpha}(M|x|)}{(M|x|)^{2-\alpha}}, \quad (4)$$

where $\langle G^2 \rangle \equiv \langle g^2(F_{\mu\nu}^a)^2 \rangle$ and $N(\alpha) > 0$ is a numerical coefficient, which will be determined. Then, in the deconfinement phase ($T > T_c$) of interest, the above chain of equalities yields

$$\text{RHS of Eq. (2)} = 2T \cdot \pi^2 2^\alpha \Gamma(\alpha) N(\alpha) \langle G^2 \rangle_T^2 M^{2\alpha-4} \sum_{k=-\infty}^{+\infty} \frac{e^{i\omega_k x_4}}{(\omega_k^2 + M^2)^\alpha},$$

where $\langle G^2 \rangle_T \equiv \langle g^2(F_{ij}^a)^2 \rangle_T$. Note that, for temperatures $T > T_*$, only the ($k = 0$)-term in the sum should be considered, since the dimensionally reduced theory is a theory of the zeroth Matsubara mode.

By using also Eq. (3), we can now rewrite Eq. (2) in terms of Fourier modes as

$$\int_0^\infty d\omega \rho(\omega) \frac{\omega}{\omega^2 + \omega_k^2} = \pi^2 2^\alpha \Gamma(\alpha) N(\alpha) \langle G^2 \rangle_T^2 \frac{M^{2\alpha-4}}{(\omega_k^2 + M^2)^\alpha}. \quad (5)$$

To solve this equation, we use the parametrization

$$\rho(\omega) = \omega f(\omega),$$

where $f(\omega)$ is some even function sufficient for the convergence of the integral at large ω 's. Motivated by earlier works [8, 10, 12], we choose it in a Lorentzian-type form

$$f(\omega) = \frac{C}{(\omega^2 + M^2)^{\alpha+\frac{1}{2}}}. \quad (6)$$

Here $C = C(T) > 0$ is some function, which will be determined. Apparently, the ω -integration in Eq. (5) converges for any choice of

$$\alpha > 0.$$

Parametrization of the spectral density through Eq. (6) guarantees furthermore that both sides of Eq. (5) have the same leading large- $|k|$ behavior. It also implies [12] that $M = M(T)$ is the momentum scale below which perturbation theory breaks down. For this reason, $M(T)$ should be of the order of the inverse correlation length of the chromo-magnetic vacuum, $\mu(T)$. Shear viscosity can finally be obtained by means of Eq. (1) as

$$\eta = \frac{\pi C}{M^{2\alpha+1}}. \quad (7)$$

We now solve Eq. (5) subsequently in the following cases: $|k| \gg 1$ and $|k| \sim 1$.

III. CONTRIBUTIONS TO η FROM HIGH AND LOW MATSUBARA MODES

A. ($|k| \gg 1$)-case

Plugging the Lorentzian-type *ansatz* (6) into Eq. (5), we obtain

$$\begin{aligned} \text{LHS of Eq. (5)} &= C \int_0^\infty d\omega \frac{\omega^2}{(\omega^2 + \omega_k^2)(\omega^2 + M^2)^{\alpha+\frac{1}{2}}} = \\ &= \frac{C}{\omega_k^{2\alpha}} \int_0^\infty dx \frac{x^2}{(x^2 + 1)(x^2 + \xi_k^2)^{\alpha+\frac{1}{2}}}, \end{aligned}$$

where $x \equiv \omega/\omega_k$ and $\xi_k \equiv M/\omega_k$. At $|k| \gg 1$, one has

$$|\xi_k| \ll 1, \quad (8)$$

and the latter integral yields

$$\text{LHS of Eq. (5)} = \frac{C}{\omega_k^{2\alpha}} \left\{ \frac{1}{\xi_k^{2\alpha}} \left[\frac{\sqrt{\pi}\Gamma(\alpha-1)}{4\Gamma(\alpha+\frac{1}{2})} \xi_k^2 + \mathcal{O}(\xi_k^3) \right] + \frac{\pi}{2\sin(\pi\alpha)} + \mathcal{O}(\xi_k^2) \right\}. \quad (9)$$

We see that, if $\alpha < 1$, then the leading term of the expansion is $\propto \frac{\pi}{2\sin(\pi\alpha)}$, i.e. it is k -independent, whereas otherwise the leading term of the expansion becomes k -dependent. For this reason, we restrict ourselves to

$$\alpha < 1$$

only. Furthermore, since $\alpha\xi_k^2 \ll 1$ as well, one can expand in powers of ξ_k also the RHS of Eq. (5) to obtain

$$\text{RHS of Eq. (5)} = \pi^2 2^\alpha \Gamma(\alpha) N(\alpha) \frac{\langle G^2 \rangle_T^2}{\omega_k^{2\alpha}} M^{2\alpha-4} [1 + \mathcal{O}(\xi_k^2)]. \quad (10)$$

Equations (9) and (10) yield the function C :

$$C = \pi 2^{\alpha+1} \Gamma(\alpha) N(\alpha) \sin(\pi\alpha) \langle G^2 \rangle_T^2 M^{2\alpha-4}.$$

Accordingly, Eq. (7) yields for the shear viscosity

$$\eta \Big|_{|k| \gg 1} = \pi^2 2^{\alpha+1} \Gamma(\alpha) N(\alpha) \sin(\pi\alpha) \frac{\langle G^2 \rangle_T^2}{M^5}. \quad (11)$$

The parametric dependence of this expression on $\langle G^2 \rangle_T$ and M is indeed the one following from the elementary dimensional analysis made in Introduction.

B. ($|k| \sim 1$)-case

Consider $|k|$'s sufficiently small for the inequality

$$|\omega_k| \ll M, \quad \text{i.e.} \quad |\xi_k| \gg 1, \quad (12)$$

to hold. Disregarding terms $\mathcal{O}(\omega_k^2/M^2)$ and higher, one has

$$\text{LHS of Eq. (5)} \simeq \frac{C}{M^{2\alpha}} \int_0^\infty \frac{dz}{(z^2 + 1)^{\alpha+\frac{1}{2}}} = \frac{\sqrt{\pi}}{2} \frac{\Gamma(\alpha)}{\Gamma(\alpha + \frac{1}{2})} \frac{C}{M^{2\alpha}},$$

where $z \equiv \omega/M$, while

$$\text{RHS of Eq. (5)} \simeq \pi^2 2^\alpha \Gamma(\alpha) N(\alpha) \frac{\langle G^2 \rangle_T^2}{M^4}.$$

We obtain from these two equations

$$C = \pi^{3/2} 2^{\alpha+1} \Gamma\left(\alpha + \frac{1}{2}\right) N(\alpha) \langle G^2 \rangle_T^2 M^{2\alpha-4}$$

and, according to Eq. (7),

$$\eta \Big|_{|k| \sim 1} = \pi^{5/2} 2^{\alpha+1} \Gamma\left(\alpha + \frac{1}{2}\right) N(\alpha) \frac{\langle G^2 \rangle_T^2}{M^5}. \quad (13)$$

In particular, at $T > T_*$, where only the ($k = 0$)-mode should be considered, this result becomes exact.




Figure 1: The ratio $\frac{\eta_{|k| \gg 1}}{\eta_{|k| \sim 1}}$ given by Eq. (25).

IV. η FROM THE CORRELATION FUNCTION $\langle T_{12}(0)T_{12}(x) \rangle$

We determine now the parameters $N(\alpha)$ and M , which enter the correlation function (4). This correlation function reads

$$\begin{aligned} \langle T_{12}(0)T_{12}(x) \rangle &= \langle g^4 F_{1\mu}^a(0)F_{2\mu}^a(0)F_{1\nu}^b(x)F_{2\nu}^b(x) \rangle = \\ &= \langle \langle g^4 F_{1\mu}^a(0)F_{2\mu}^a(0)F_{1\nu}^b(x)F_{2\nu}^b(x) \rangle \rangle + \langle g^2 F_{1\mu}^a(0)F_{2\mu}^a(0) \rangle \langle g^2 F_{1\nu}^b(x)F_{2\nu}^b(x) \rangle + \\ &+ \langle g^2 F_{1\mu}^a(0)F_{1\nu}^b(x) \rangle \langle g^2 F_{2\mu}^a(0)F_{2\nu}^b(x) \rangle + \langle g^2 F_{1\mu}^a(0)F_{2\nu}^b(x) \rangle \langle g^2 F_{2\mu}^a(0)F_{1\nu}^b(x) \rangle, \end{aligned} \quad (14)$$

where double angular brackets denote a connected (or irreducible) average. We use the Gaussian-dominance hypothesis [11], which allows one to disregard this connected average. For the two-point correlation function of gluonic field strengths we use the standard parametrization [11, 19]

$$\langle g^2 F_{\mu\nu}^a(x)F_{\lambda\rho}^b(x') \rangle = (\delta_{\mu\lambda}\delta_{\nu\rho} - \delta_{\mu\rho}\delta_{\nu\lambda}) \frac{\langle G^2 \rangle}{12(N_c^2 - 1)} \delta^{ab} D(x - x'), \quad (15)$$

where $D(x)$ is a dimensionless function mediating the confining interaction. In this parametrization, we have disregarded a small contribution of non-confining non-perturbative interactions [17], [26]. By using Eq. (15), we obtain for the correlation function (14):

$$\langle T_{12}(0)T_{12}(x) \rangle \simeq \langle g^2 F_{1\mu}^a(0)F_{2\mu}^a(0) \rangle \langle g^2 F_{1\nu}^b(x)F_{2\nu}^b(x) \rangle + \langle g^2 F_{1\mu}^a(0)F_{1\nu}^b(x) \rangle \langle g^2 F_{2\mu}^a(0)F_{2\nu}^b(x) \rangle +$$

$$+ \langle g^2 F_{1\mu}^a(0) F_{2\nu}^b(x) \rangle \langle g^2 F_{2\mu}^a(0) F_{1\nu}^b(x) \rangle = \frac{\langle G^2 \rangle^2}{72(N_c^2 - 1)} D^2(x). \quad (16)$$

The dimensionless function $D(x)$ is usually chosen in the form

$$D(x) = e^{-\mu|x|}. \quad (17)$$

Inserting this expression into the formula for the string tension in the fundamental representation,

$$\sigma_f = \frac{\langle G^2 \rangle}{144} \int d^2x D(x), \quad (18)$$

one can define the gluon condensate in terms of σ_f and the vacuum correlation length μ as follows [17, 19]:

$$\langle G^2 \rangle = \frac{72}{\pi} \sigma_f \mu^2. \quad (19)$$

To obtain for the correlator $\langle T_{12}(0) T_{12}(x) \rangle$ the functional form given by the RHS of Eq. (4), we modify parametrization (17) to

$$D(x) = \mathcal{A}(\alpha) \sqrt{\frac{K_{2-\alpha}(2\mu|x|)}{(2\mu|x|)^{2-\alpha}}}, \quad (20)$$

where $\mathcal{A}(\alpha)$ is a numerical normalization factor. At $|x| \gtrsim \mu^{-1}$, the new function (20) falls off with the same exponent as Eq. (17). To obtain the normalization factor $\mathcal{A}(\alpha)$, we substitute Eq. (20) into relation (18), which holds for any function $D(x)$. Using further expression (19), we obtain

$$\mathcal{A}(\alpha) = \frac{4}{\int_0^\infty dz \sqrt{z^\alpha K_{2-\alpha}(z)}}. \quad (21)$$

The correlator (16) now reads

$$\langle T_{12}(0) T_{12}(x) \rangle = \frac{\mathcal{A}^2(\alpha)}{576} \langle G^2 \rangle^2 \frac{K_{2-\alpha}(2\mu|x|)}{(2\mu|x|)^{2-\alpha}}, \quad (22)$$

where the function $\mathcal{A}(\alpha)$ is given by Eq. (21), and we have fixed $N_c = 3$. Comparing Eq. (22) with the original definition (4), we conclude that

$$N(\alpha) = \frac{\mathcal{A}^2(\alpha)}{576} \quad \text{and} \quad M = 2\mu.$$

Equations (11) and (13) yield now contributions to η from high and low Matsubara modes:

$$\eta \Big|_{|k| \gg 1} = \frac{\pi^2}{9216} 2^\alpha \Gamma(\alpha) \sin(\pi\alpha) \frac{[\mathcal{A}(\alpha) \langle G^2 \rangle_T]^2}{\mu^5(T)} \quad (23)$$




Figure 2: Entropy density $s(T)$ in the units of T^3 obtained from the lattice values for the pressure p_{lat} [16] (courtesy of F. Karsch).

and

$$\eta\Big|_{|k|\sim 1} = \frac{\pi^{5/2}}{9216} 2^\alpha \Gamma\left(\alpha + \frac{1}{2}\right) \frac{[\mathcal{A}(\alpha) \langle G^2 \rangle_T]^2}{\mu^5(T)}. \quad (24)$$

The ratio of these results,

$$\frac{\eta\Big|_{|k|\gg 1}}{\eta\Big|_{|k|\sim 1}} = \frac{\Gamma(\alpha) \sin(\pi\alpha)}{\sqrt{\pi} \Gamma\left(\alpha + \frac{1}{2}\right)}, \quad (25)$$

in the interval $0 < \alpha < 1$ of interest is plotted in Fig. 1. It equals to unity at $\alpha = 1/2$, i.e., at this value of α , our results for shear viscosity become k -independent, as they should be.

This yields the principal analytic result of the present paper:

$$\eta(T) = \frac{\pi^{5/2}}{4608\sqrt{2}} \frac{[\mathcal{A}(1/2) \langle G^2 \rangle_T]^2}{\mu^5(T)}, \quad (26)$$

where $\mathcal{A}(1/2) \simeq 1.05$. Remarkably, *ansatz* (6) at $\alpha = 1/2$ takes the conventional Lorentzian form. In the next Section, we will evaluate the ratio η/s numerically.

V. NUMERICAL EVALUATION

Following Ref. [16], we assume the value $T_c = 270 \text{ MeV}$ in $\text{SU}(3)$ YM theory. We use the two-loop running coupling [16]

$$g^{-2}(T) = 2b_0 \ln \frac{T}{\Lambda} + \frac{b_1}{b_0} \ln \left(2 \ln \frac{T}{\Lambda} \right), \text{ where } b_0 = \frac{11N_c}{48\pi^2}, \quad b_1 = \frac{34}{3} \left(\frac{N_c}{16\pi^2} \right)^2, \quad \Lambda = 0.104T_c,$$

and $N_c = 3$ for the case under study. We also assume for $\mu(T)$ and for the spatial string tension in the fundamental representation, $\sigma_f(T)$, the following parametrizations [15, 17]:

$$\mu(T) = \mu \cdot \begin{cases} 1 & \text{at } T_c < T < T_*, \\ \frac{g^2(T)T}{g^2(T_{\text{d.r.}})T_{\text{d.r.}}} & \text{at } T > T_*, \end{cases} \quad (27)$$

$$\sigma_f(T) = \sigma_f \cdot \begin{cases} 1 & \text{at } T_c < T < T_*, \\ \left[\frac{g^2(T)T}{g^2(T_{\text{d.r.}})T_{\text{d.r.}}} \right]^2 & \text{at } T > T_*, \end{cases} \quad (28)$$

where $\mu = 894 \text{ MeV}$ [13] and $\sigma_f = (0.44 \text{ GeV})^2$. Equation (19), extrapolated to finite temperatures, yields for the chromo-magnetic gluon condensate $\langle G^2 \rangle_T$ [15, 17]:

$$\langle G^2 \rangle_T = \frac{72}{\pi} \sigma_f(T) \mu^2(T) = \langle G^2 \rangle \cdot \begin{cases} 1 & \text{at } T_c < T < T_*, \\ \left[\frac{g^2(T)T}{g^2(T_{\text{d.r.}})T_{\text{d.r.}}} \right]^4 & \text{at } T > T_*. \end{cases}$$

The value of T_* can be estimated from the equation

$$\sigma_f(T_*) = \sigma_f,$$

where $\sigma_f(T) = [0.566g^2(T)T]^2$ is the high-temperature parametrization of the fundamental spatial string tension [16]. Solving this equation numerically, one obtains [20]

$$T_* = 1.28T_c.$$

The entropy density $s = s(T)$ can be obtained by the formula $s = \partial p_{\text{lat}} / \partial T$, where we use for the pressure p_{lat} the corresponding lattice values from Ref. [16]. In Fig. 2, we plot $s(T)$ in the units of T^3 . The temperature dependence of the ratio η/s is determined by the function $\langle G^2 \rangle_T^2 / [\mu^5(T)s(T)]$. One can check numerically that, at $T \gtrsim 2T_c$ where s/T^3 is nearly constant, $\langle G^2 \rangle_T^2 / [\mu^5(T)s(T)] = \mathcal{O}(g^6(T))$, as was mentioned in Introduction.

In Fig. 3, we plot the ratio η/s , with η given by Eq. (26), as a function of temperature. Also in Fig. 3, we plot the conjectured lower bound for this ratio, equal to $(4\pi)^{-1} \simeq 0.08$,




Figure 3: Calculated values of the ratio η/s as a function of temperature. Also shown is the conjectured lower bound of $(4\pi)^{-1}$ for this quantity, realized in $\mathcal{N} = 4$ SYM.

which is realized in $\mathcal{N} = 4$ SYM [5]. This bound is indeed not reached by our values, although they get very close to it at the highest temperature $T = 4.54T_c$ where the lattice data for the pressure (and therefore also for s) are available.

Furthermore, we compare numerically the obtained nonperturbative spectral density,

$$\rho(\omega, T) = C(T) \frac{\omega}{\omega^2 + 4\mu^2(T)}, \quad \text{where } C(T) = \left(\frac{\pi}{2}\right)^{3/2} \frac{\mathcal{A}^2(1/2)}{576} \frac{\langle G^2 \rangle_T^2}{\mu^3(T)}, \quad (29)$$

with the perturbative one, which at the tree level reads [9, 10]

$$\rho^{\text{pert}}(\omega, T) = \frac{1}{20\pi^2} \theta(\omega - \tilde{\omega}(T)) \frac{\omega^4}{\tanh \frac{\omega}{4T}}.$$

Note that $\rho^{\text{pert}}(\omega, T) \neq 0$ only at $\omega > \tilde{\omega}(T)$, where $\tilde{\omega}(T) \simeq 7.5T$ [10]. For this reason, $\rho^{\text{pert}}(\omega, T)$ in any case does not affect the calculated η , which is defined according to Eq. (1) by the values of the spectral density at $\omega \rightarrow 0$. Figure 4 illustrates the full spectral density $\rho^{\text{full}} = \rho + \rho^{\text{pert}}$ as a function of ω/T at $T = T_c$. At $\omega < \tilde{\omega}(T)$, ρ^{full} is given by the obtained result (29), while at $\omega > \tilde{\omega}(T)$ the perturbative part ρ^{pert} takes it over. Were ρ^{pert} nonvanishing down to $\omega = 0$, it would dominate over ρ already at $\omega > 4.85T$. That is the reason why, by $\omega = \tilde{\omega}(T)$, ρ^{pert} significantly exceeds ρ , as one can see from the gap in the

values of ρ^{full} at this value of ω . Qualitatively, the same ρ^{full} and the relation between ρ and ρ^{pert} persist with the increase of temperature.

Finally, in Appendix A, we illustrate the correspondence between the splitting of $\rho^{\text{full}} = \rho + \rho^{\text{pert}}$ and the splitting of $\langle T_{12}(0)T_{12}(x) \rangle_{\text{full}} = \langle T_{12}(0)T_{12}(x) \rangle + \langle T_{12}(0)T_{12}(x) \rangle_{\text{pert}}$. This correspondence allows one to isolate the contribution, which ρ^{pert} brings about to the ω -integral in the full Kubo formula.

VI. DISCUSSION AND OUTLOOK

In this paper, we have applied Kubo formula to a nonperturbative calculation of the shear viscosity η in SU(3) YM theory. With the use of the stochastic vacuum model, the $\langle T_{12}(0)T_{12}(x) \rangle$ -correlator entering Kubo formula has been expressed in terms of the temperature-dependent chromo-magnetic gluon condensate $\langle g^2(F_{ij}^a)^2 \rangle_T$ and the correlation length of the chromo-magnetic vacuum $\mu^{-1}(T)$. As was expected (cf. Introduction), η turns out to be $\propto \mu^{-5}(T) \langle g^2(F_{ij}^a)^2 \rangle_T^2$, where the numerical factor is given by Eq. (26). At temperatures $T \gtrsim 2T_c$, the calculated ratio η/s falls off as $\mathcal{O}(g^6(T))$, as it should do in the dimensionally-reduced theory. Numerically, up to the temperature $T = 4.54T_c$, where the lattice data on bulk thermodynamic quantities are still available, the obtained values of the ratio η/s stay above the conjectured lower bound of $(4\pi)^{-1}$, which is reached in $\mathcal{N} = 4$ SYM.

Formally, our result (26) persists even at higher temperatures, being extrapolated to which it yields for the (η/s) -ratio values smaller than $(4\pi)^{-1}$. One should, however, realize that the monotonic fall-off of η/s with temperature, stemming from the relation $\eta \propto \mu^{-5}(T) \langle g^2(F_{ij}^a)^2 \rangle_T^2$, is predefined by our calculational method, which combines Kubo formula with the stochastic vacuum model. In fact, all the kinetic coefficients derivable in this way should be $\propto \langle g^2(F_{ij}^a)^2 \rangle_T^2$ (cf. Introduction). In particular, this is the case for the bulk viscosity ζ [12], which can be obtained from the correlation function

$$\langle g^4 F_{\mu\nu}^{a2}(0) F_{\lambda\rho}^{b2}(x) \rangle \simeq \langle G^2 \rangle^2 \left[1 + \frac{1}{3(N_c^2 - 1)} D^2(x) \right].$$

(Here “ $\simeq$ ” stands for “Gaussian approximation”.) On general grounds [23], one indeed expects a monotonic fall-off of the (ζ/s) -ratio with temperature, as was confirmed by explicit calculations [12]. However, on the same general grounds [23], for the (η/s) -ratio in question




Figure 4: The full spectral density $\rho^{\text{full}}(\omega, T)$ as a function of ω/T at $T = T_c$. At $\frac{\omega}{T} < 7.5$, ρ^{full} is entirely nonperturbative and given by Eq. (29), while at $\frac{\omega}{T} > 7.5$ the perturbative part ρ^{pert} is dominating.

one expects the existence of a minimum at temperatures close to T_c [27] and a subsequent *increase* with the further increase of temperature. Indeed, at least at the temperature as high as $47.4T_c$, $g(T)$ becomes smaller than unity, and the weakly-interacting dilute-gas model of the gluon plasma gradually sets in. As mentioned in Introduction, in the dilute-gas model [25] $(\eta/s) \sim 1/(g^4 \ln g^{-1})$, i.e. this ratio increases with temperature. The stochastic vacuum model, on the other hand, being applicable at strong coupling, correctly yields the expected fall-off of the (η/s) -ratio at temperatures $\lesssim 2T_c$, but cannot reproduce its increase at much higher temperatures.

We would also like to emphasize an interesting fact, which has been realized by the end of the calculation. We have started with the general α -dependent Lorentzian-type *ansatz* (6) for the spectral density $\rho(\omega)$. By using it in the Kubo formula, we have come to the conclusion that only for the single value, $\alpha = 1/2$, this *ansatz* provides the Matsubara-mode independence of $\rho(\omega)$. For this value of α , Eq. (6) takes the conventional Lorentzian form. In this way, also the function $D(x)$ from Eq. (15) is defined unambiguously as

$$D(x) = \mathcal{A}(1/2) \sqrt{\frac{K_{3/2}(2\mu|x|)}{(2\mu|x|)^{3/2}}},$$

where the value of $\mathcal{A}(1/2)$ can be found after Eq. (26).

Note finally that we have used in our calculation the Gaussian-dominance hypothesis [11], which allows one to disregard in Eq. (14) the connected four-point correlation function of gluonic field strengths compared to the pairwise products of the two-point correlation functions. The same approximation was used in Ref. [21] for the calculation of topological susceptibility via the four-point correlation function. This approximation can be relaxed in the following way. Consider a parametrization for the nonperturbative part of the connected four-point correlation function suggested in Ref. [22]:

$$\begin{aligned}
& \langle \langle g^4 F_{\mu_1 \nu_1}^{a_1}(x_1) F_{\mu_2 \nu_2}^{a_2}(x_2) F_{\mu_3 \nu_3}^{a_3}(x_3) F_{\mu_4 \nu_4}^{a_4}(x_4) \rangle \rangle = \\
& = \langle G^2 \rangle^2 \left\{ f^{a_1 a_2 b} f^{a_3 a_4 b} (\varepsilon_{\mu_1 \nu_1 \mu_3 \nu_3} \varepsilon_{\mu_2 \nu_2 \mu_4 \nu_4} - \varepsilon_{\mu_1 \nu_1 \mu_4 \nu_4} \varepsilon_{\mu_2 \nu_2 \mu_3 \nu_3}) + \right. \\
& \quad + f^{a_3 a_1 b} f^{a_2 a_4 b} (\varepsilon_{\mu_1 \nu_1 \mu_4 \nu_4} \varepsilon_{\mu_2 \nu_2 \mu_3 \nu_3} - \varepsilon_{\mu_1 \nu_1 \mu_2 \nu_2} \varepsilon_{\mu_3 \nu_3 \mu_4 \nu_4}) + \\
& \quad + f^{a_2 a_3 b} f^{a_1 a_4 b} (\varepsilon_{\mu_1 \nu_1 \mu_2 \nu_2} \varepsilon_{\mu_3 \nu_3 \mu_4 \nu_4} - \varepsilon_{\mu_1 \nu_1 \mu_3 \nu_3} \varepsilon_{\mu_2 \nu_2 \mu_4 \nu_4}) - \\
& \quad - \frac{1}{8} \left[\delta^{a_1 a_2} \delta^{a_3 a_4} (\delta_{\mu_1 \mu_2} \delta_{\nu_1 \nu_2} - \delta_{\mu_1 \nu_2} \delta_{\mu_2 \nu_1}) (\delta_{\mu_3 \mu_4} \delta_{\nu_3 \nu_4} - \delta_{\mu_3 \nu_4} \delta_{\mu_4 \nu_3}) + \right. \\
& \quad + \delta^{a_1 a_3} \delta^{a_2 a_4} (\delta_{\mu_1 \mu_3} \delta_{\nu_1 \nu_3} - \delta_{\mu_1 \nu_3} \delta_{\mu_3 \nu_1}) (\delta_{\mu_2 \mu_4} \delta_{\nu_2 \nu_4} - \delta_{\mu_2 \nu_4} \delta_{\mu_4 \nu_2}) + \\
& \quad \left. + \delta^{a_1 a_4} \delta^{a_2 a_3} (\delta_{\mu_1 \mu_4} \delta_{\nu_1 \nu_4} - \delta_{\mu_1 \nu_4} \delta_{\mu_4 \nu_1}) (\delta_{\mu_2 \mu_3} \delta_{\nu_2 \nu_3} - \delta_{\mu_2 \nu_3} \delta_{\mu_3 \nu_2}) \right] \Big\} \tilde{D}(z_1, \dots, z_6), \quad (30)
\end{aligned}$$

where $z_1 = x_1 - x_2$, $z_2 = x_1 - x_3, \dots$, $z_6 = x_3 - x_4$ are relative coordinates. For the connected correlation function entering Eq. (14) this parametrization yields

$$\langle \langle g^4 F_{1\mu}^a(0) F_{2\mu}^a(0) F_{1\nu}^b(x) F_{2\nu}^b(x) \rangle \rangle = 2(N_c^2 - 1) \left(N_c - \frac{1}{8} \right) \langle G^2 \rangle^2 \tilde{D}(0, x, x, x, x, 0).$$

Similarly to Eq. (20), for the function $\tilde{D}$ too one can have a parametrization compatible with Eq. (4):

$$\tilde{D}(0, x, x, x, x, 0) = \mathcal{B}(\alpha) \frac{K_{2-\alpha}(4\mu|x|)}{(4\mu|x|)^{2-\alpha}}.$$

The normalization factor $\mathcal{B}(\alpha)$ should now be determined simultaneously with the normalization factor $\mathcal{A}(\alpha)$ from a system of equations for two observables, which both depend on the functions D and $\tilde{D}$. Natural observables of this kind are the string tension and the topological susceptibility. The contribution of the function $\tilde{D}$ to the string tension has already been evaluated in Ref. [22]. Further analysis of the outlined extension of the Gaussian approximation is, however, not the purpose of the present paper.

Acknowledgments

I am grateful to Frithjof Karsch, Hans-Jürgen Pirner and Arif Shoshi for helpful discussions, and to Yoshimasa Hidaka and Olaf Kaczmarek for the useful correspondence and comments. I also thank Frithjof Karsch for providing the details of the lattice data from Ref. [16]. This work has been supported by the German Research Foundation (DFG), contract Sh 92/2-1.

Appendix A. Matching perturbative contributions in the Kubo formula.

For this Appendix, we promote Eq. (2) to the full Kubo formula, i.e. replace ρ by $\rho^{\text{full}} = \rho + \rho^{\text{pert}}$ and $\langle T_{12}(0)T_{12}(x) \rangle$ by $\langle T_{12}(0)T_{12}(x) \rangle_{\text{full}} = \langle T_{12}(0)T_{12}(x) \rangle + \langle T_{12}(0)T_{12}(x) \rangle_{\text{pert}}$. On the LHS of such a full Kubo formula, consider the integral containing ρ_{pert} . To facilitate the ω -integration, we approximate the perturbative part of the spectral density by the function $\rho^{\text{pert}} = N\omega^4$ down to $\omega = 0$ and determine the coefficient N . The integral emerging on the LHS of the Kubo formula then reads

$$\begin{aligned} \int_0^\infty d\omega \omega^4 [\cosh(\omega x_4) \coth(\omega\beta/2) - \sinh(\omega x_4)] &= \sum_{n=0}^\infty \left[\frac{1}{(\beta n + x_4)^5} + \frac{1}{[\beta n + (\beta - x_4)]^5} \right] = \\ &= 24 \left\{ \frac{1}{x_4^5} + \sum_{n=1}^\infty \left[\frac{1}{(\beta n + x_4)^5} + \frac{1}{(\beta n - x_4)^5} \right] \right\}. \end{aligned} \quad (\text{A.1})$$

To obtain the last expression, we have extracted the $(n = 0)$ -term from the sum $\sum_{n=0}^\infty \frac{1}{(\beta n + x_4)^5}$ and shifted n by 1 in the sum $\sum_{n=0}^\infty \frac{1}{[\beta n + (\beta - x_4)]^5}$. Note that $x_4 \in [0, \beta]$, and the obtained expression has singularities at $x_4 = 0$ and $x_4 = \beta$ [in the $(n = 1)$ -term]. These singularities are identical, since $\cosh[\omega(x_4 - \frac{\beta}{2})] = \cosh(\omega\beta/2)$ both at $x_4 = 0$ and $x_4 = \beta$.

We will demonstrate now that this expression corresponds to the contribution, which $\langle T_{12}(0)T_{12}(x) \rangle_{\text{pert}}$ brings about to the RHS of the full Kubo formula. The UV-finite part of this perturbative correlation function can be written as

$$\langle T_{12}(0)T_{12}(x) \rangle_{\text{pert}} = \frac{A(N_c^2 - 1)g^4}{x^8},$$

where the value of the numerical constant A depends on the regularization scheme applied. Thus, the sum emerging on the RHS of the Kubo formula reads

$$\sum_{n=-\infty}^{+\infty} \frac{1}{[\mathbf{x}^2 + (\beta n + x_4)^2]^4}.$$

Doing the integration over d^3x first, we have

$$\int_0^\infty dx \frac{x^2}{[x^2 + (\beta n + x_4)^2]^4} = \frac{\pi}{32|\beta n + x_4|^5}.$$

The part of the sum with negative winding modes reads

$$\sum_{n=-\infty}^{-1} \frac{1}{|\beta n + x_4|^5} = \sum_{n=-\infty}^{-1} \frac{1}{(-\beta n - x_4)^5} = \sum_{n=1}^{\infty} \frac{1}{(\beta n - x_4)^5}.$$

Therefore, the integration over $d|\mathbf{x}|$ and the summation over winding modes yield

$$\int_0^\infty dx x^2 \sum_{n=-\infty}^{+\infty} \frac{1}{[x^2 + (\beta n + x_4)^2]^4} = \frac{\pi}{32} \left\{ \frac{1}{x_4^5} + \sum_{n=1}^{\infty} \left[\frac{1}{(\beta n + x_4)^5} + \frac{1}{(\beta n - x_4)^5} \right] \right\}. \quad (\text{A.2})$$

The coincidence of curly brackets in Eqs. (A.1) and (A.2) proves that the *ansatz* $\rho^{\text{pert}} = N\omega^4$ captures the contribution of $\langle T_{12}(0)T_{12}(x) \rangle_{\text{pert}}$ correctly, with the corresponding normalization factor being $N(T) = \pi^2 A(T)(N_c^2 - 1)g^4(T)/192$.

In this way, one isolates in the full Kubo formula simultaneously the perturbative ω^4 -part of ρ^{full} and the perturbative contribution to $\langle T_{12}(0)T_{12}(x) \rangle_{\text{full}}$. The remaining nonperturbative parts of ρ^{full} and $\langle T_{12}(0)T_{12}(x) \rangle_{\text{full}}$ are related to each other by means of Eq. (2).

-
- [1] For a review see e.g.: K. Yagi, T. Hatsuda and Y. Miake, *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **23**, 1 (2005).
 - [2] H. Appelshäuser *et al.* [NA49 Collaboration], *Phys. Rev. Lett.* **80**, 4136 (1998); A. M. Poskanzer *et al.* [NA49 Collaboration], *Nucl. Phys. A* **661**, 341 (1999); K. H. Ackermann *et al.* [STAR Collaboration], *Phys. Rev. Lett.* **86**, 402 (2001).
 - [3] K. Adcox *et al.* [PHENIX Collaboration], *Nucl. Phys. A* **757**, 184 (2005); I. Arsene *et al.* [BRAHMS Collaboration], *Nucl. Phys. A* **757**, 1 (2005); B. B. Back *et al.* [PHOBOS Collaboration], *Nucl. Phys. A* **757**, 28 (2005); J. Adams *et al.* [STAR Collaboration], *Nucl. Phys. A* **757**, 102 (2005).
 - [4] D. Molnar and M. Gyulassy, *Nucl. Phys. A* **697**, 495 (2002) [Erratum-ibid. A **703**, 893 (2002)].
 - [5] G. Policastro, D. T. Son and A. O. Starinets, *Phys. Rev. Lett.* **87**, 081601 (2001).
 - [6] Z. Xu and C. Greiner, *Phys. Rev. Lett.* **100**, 172301 (2008); Z. Xu, C. Greiner and H. Stöcker, *Phys. Rev. Lett.* **101**, 082302 (2008).

- [7] M. Asakawa, S. A. Bass and B. Müller, Phys. Rev. Lett. **96**, 252301 (2006); Prog. Theor. Phys. **116**, 725 (2007).
- [8] F. Karsch and H. W. Wyld, Phys. Rev. D **35**, 2518 (1987).
- [9] G. Aarts and J. M. Martinez Resco, JHEP **04**, 053 (2002).
- [10] H. B. Meyer, Phys. Rev. D **76**, 101701 (2007).
- [11] For reviews see: A. Di Giacomo, H. G. Dosch, V. I. Shevchenko and Yu. A. Simonov, Phys. Rept. **372**, 319 (2002); D. Antonov, Surv. High Energ. Phys. **14**, 265 (2000).
- [12] D. Kharzeev and K. Tuchin, JHEP **09**, 093 (2008); F. Karsch, D. Kharzeev and K. Tuchin, Phys. Lett. B **663**, 217 (2008).
- [13] A. Di Giacomo, E. Meggiolaro and H. Panagopoulos, Nucl. Phys. B **483**, 371 (1997); M. D’Elia, A. Di Giacomo and E. Meggiolaro, Phys. Rev. D **67**, 114504 (2003).
- [14] G. S. Bali, N. Brambilla and A. Vairo, Phys. Lett. B **421**, 265 (1998).
- [15] N. O. Agasian, Phys. Lett. B **562**, 257 (2003).
- [16] G. Boyd, J. Engels, F. Karsch, E. Lärermann, C. Legeland, M. Lütgemeier and B. Petersson, Nucl. Phys. B **469**, 419 (1996).
- [17] D. Antonov and H. J. Pirner, Eur. Phys. J. C **55**, 439 (2008).
- [18] O. Nachtmann, Annals Phys. **209**, 436 (1991); S. Donnachie, H. G. Dosch, O. Nachtmann and P. Landshoff, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **19**, 1 (2002).
- [19] A. I. Shoshi, F. D. Steffen, H. G. Dosch and H. J. Pirner, Phys. Rev. D **68**, 074004 (2003).
- [20] D. Antonov, H. J. Pirner and M. G. Schmidt, “Interaction measure $(\varepsilon - 3p)/T^4$ of the gluon plasma,” arXiv:0808.2201 [hep-ph].
- [21] M. N. Chernodub and I. E. Kozlov, JETP Lett. **86**, 1 (2007).
- [22] W. Kornelis and H. G. Dosch, Nucl. Phys. Proc. Suppl. **96**, 426 (2001).
- [23] L. P. Csernai, J. I. Kapusta and L. D. McLerran, Phys. Rev. Lett. **97**, 152303 (2006); J. I. Kapusta, “Viscous properties of strongly interacting matter at high temperature,” arXiv:0809.3746 [nucl-th].
- [24] Y. Hidaka and R. D. Pisarski, Phys. Rev. D **78**, 071501 (2008).
- [25] P. Arnold, G. D. Moore and L. G. Yaffe, JHEP **11**, 001 (2000); *ibid.* **05**, 051 (2003).
- [26] High-energy scattering data in the vacuum suggest that the relative weight of these interactions constitutes of at most 26% [19].
- [27] Such a minimum appears, for example, upon the multiplication of the perturbative result for

the (η/s) -ratio [25] by the squared fundamental Polyakov loop [24].